

An improved calibration for the mixing length theory of convection

Federico Spada




Max-Planck-Institut für
Sonnensystemforschung

In collaboration with: P. Demarque, J.D. Tanner, S. Basu (Yale)

Convection in stars



> 1.5 solar masses



0.5 - 1.5 solar masses




< 0.5 solar masses



Low-mass stars have convectively unstable outer layers

Outer CZs affect overall structure




Mixing Length Theory (MLT)

- MLT contains free parameter α
- Standard calibration of α based on the Sun (t_{\odot} , M_{\odot} , R_{\odot})



- Solar-calibrated α :
 - constant along evolutionary track
 - independent of other stellar parameters (e.g., mass, evolutionary phase, metallicity)

very strong assumptions!

$$\alpha = \ell / H_P$$




Radius inflation of low-mass stars



- $M < 0.7 M_{\odot}$ stars have radii larger than model predictions by $\sim 3\text{-}5\%$
- Rotational/magnetic effects on convection are a natural explanation


3D RHD simulations of CZs



Ludwig et al., A&A (1999)

s_{ad} : adiabatic specific entropy (units: $\text{erg g}^{-1} \text{K}^{-1}$) in the deep convective layers

Calibration of α using RHD simulations



Tune α to reproduce the value of s_{ad} obtained from RHD simulations

Extract s_{ad} from 3D RHD simulations


One-dimensional fit:

$$s_{\text{ad}}^{\text{RHD}} = s_0 + \beta \exp\left(\frac{x - x_0}{\tau}\right)$$


as a function of the variable:

$$x = A \log T_{\text{eff}} + B \log g$$


The parameters of the fit are metallicity-dependent



Entropy-based calibration of α



$\rightarrow \alpha_{\text{MLT}}^{\text{cal}}([\text{Fe}/\text{H}], x);$



Spada, Demarque, Basu, and Tanner (in preparation)

Implementation in a stellar code (YREC)

Construction of an entropy-calibrated evolutionary track:

At each evolutionary step:



- From the current stellar parameters, evaluate:


$$s_{\text{ad}}^{\text{RHD}} = s_0 + \beta \exp\left(\frac{x - x_0}{\tau}\right)$$

- Iteratively solve (e.g., bisection method):


$$s_{\text{ad}}^{\text{st. code}}(\alpha_{\text{MLT}}) \equiv s_{\text{ad}}^{\text{RHD}} \rightarrow \alpha_{\text{MLT}}^{\text{cal}}([\text{Fe}/\text{H}], x)$$

1 M_{\odot} evolutionary track







**Luminosity
is unaffected**




**Significant impact on radius,
properties of the CZ (e.g., depth,
bottom temperature, S_{ad})**



Impact on light elements depletion

**Li destruction during the PMS
is affected (~factor of 2)**



Final Remarks

- Entropy-based α yields an improved calibration for the radii of low-mass stars (radius discrepancy problem)
- More physically-motivated than solar-calibration: comparing α across stellar evolution codes is meaningless
- Naturally takes into account effect on convection of stellar parameters (mass, surface composition, evolutionary phase)
- Yields improved evolutionary history of the Sun (potential impact on dynamics, habitability of solar system objects)